


## TdT Antibody / DNA nucleotidylexotransferase (R31581)

Catalog No.	Formulation	Size
R31581	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
<b>Gene ID</b>	1791
<b>Applications</b>	Western Blot : 0.5-1ug/ml
<b>Limitations</b>	This TdT antibody is available for research use only.



Western blot testing of TdT antibody and Jurkat lysate. Predicted molecular weight: ~56 kDa.



Western blot testing of TdT antibody and recombinant human protein (0.5ng)

## Description

Terminal Deoxynucleotidyl Transferase, also known as Terminal transferase, is a unique DNA polymerase that without template direction catalyzes the addition of deoxyribonucleotides onto the 3-prime-hydroxyl end of DNA primers. Its gene is mapped to the region 10q23-q24. And TdT cDNA contains an open reading frame of 1,530 basepairs corresponding to a protein containing 510 amino acids. TdT may be responsible for inserting nucleotides (N regions) at the V(H)-D and D-J(H) junctions of immunoglobulin genes. The enzyme is present in immature thymocytes, some bone marrow cells, transformed pre-B and pre-T cell lines, and leukemia cells. Additionally, TdT catalyses the addition of nucleotides to the 3' terminus of a DNA molecule. Unlike most DNA polymerases it does not require a template. The preferred substrate of this enzyme is a 3'-overhang, but it can also add nucleotides to blunt or recessed 3' ends. Cobalt is a necessary cofactor.

## Application Notes

The stated application concentrations are suggested starting amounts. Titration of the TdT antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

Human partial recombinant protein (AA 316-509) was used as the immunogen for this TdT antibody.

## Storage

After reconstitution, the TdT antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.