


# **THBS1 Antibody / Thrombospondin (RQ5842)**


Catalog No.	Formulation	Size
RQ5842	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P07996
Applications	Western Blot : 0.5-1ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This THBS1 antibody is available for research use only.



Flow cytometry testing of human U-2 OS cells with THBS1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= THBS1 antibody.



Western blot testing of human Jurkat cell lysate with THBS1 antibody. Expected molecular weight: 130-200 kDa depending on glycosylation level.

### **Description**

THBS1 is also known as Thrombospondin 1, TSP1. The protein encoded by this gene is a subunit of a disulfide-linked homotrimeric protein. It is an adhesive glycoprotein that mediates cell-to-cell and cell-to-matrix interactions. Also this protein can bind to fibrinogen, fibronectin, laminin, type V collagen and integrins alpha-V/beta-1. This protein has been shown to play roles in platelet aggregation, angiogenesis, and tumorigenesis. In addition, the thrombospondin-1 protein is a member of the thrombospondinfamily. It is a multi-domain matrix glycoprotein that has been shown to be a natural inhibitor of neovascularization and tumorigenesis in healthy tissue. Both positive and negative modulation of endothelial cell adhesion, motility, and growth have been attributed to TSP1. This should not be surprising considering that TSP1 interacts with at least 12 cell adhesion receptors, including CD36, ?v integrins, ?1 integrins, syndecan, and integrinassociated protein (IAP or CD47). It also interacts with numerous proteases involved in angiogenesis, including plasminogen, urokinase, matrix metalloproteinase, thrombin, cathepsin, and elastase.

### **Application Notes**

Optimal dilution of the THBS1 antibody should be determined by the researcher.

### **Immunogen**

Recombinant human protein (amino acids N19-A268) was used as the immunogen for the THBS1 antibody.

#### **Storage**

After reconstitution, the THBS1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.