


TP53I13 Antibody / Tumor protein p53-inducible protein 13 (RQ7852)


Catalog No.	Formulation	Size
RQ7852	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q8NBR0
Localization	Cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence : 5ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This TP53I13 antibody is available for research use only.




IHC staining of FFPE human spleen tissue with TP53I13 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human endometrial adenocarcinoma tissue with TP53I13 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human colon cancer tissue with TP53I13 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human ovarian cancer tissue with TP53I13 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human liver cancer tissue with TP53I13 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human placental tissue with TP53I13 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human placental tissue with TP53I13 antibody (red) and DAPI nuclear stain (blue). HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human HaCaT, 2) mouse heart and 3) rat heart tissue lysate with TP53I13 antibody. Predicted molecular weight ~42 kDa.

Description

Involved in several processes, including negative regulation of cell cycle; response to UV; and response to xenobiotic stimulus. Located in cytoplasm.

Application Notes

Optimal dilution of the TP53I13 antibody should be determined by the researcher.

Immunogen

E. coli-derived recombinant human protein (amino acids A30-Q365) was used as the immunogen for the TP53I13 antibody.

Storage

After reconstitution, the TP53I13 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.