


TRAF1 Antibody (R34527)

| Catalog No. | Formulation | Size |
|--------------|---|--------|
| R34527-100UG | 0.5~mg/ml in 1X TBS, pH7.3, with 0.5% BSA (US sourced) and 0.02% sodium azide | 100 ug |


Bulk quote request

| Availability | 1-3 business days |
|----------------------|---|
| Species Reactivity | Human |
| Predicted Reactivity | Mouse |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (goat origin) |
| Isotype | Goat Ig |
| Purity | Antigen affinity |
| Gene ID | 7185 |
| Localization | Cytoplasmic |
| Applications | Western Blot : 0.5-1.5ug/ml IHC (FFPE) : 2.5ug/ml ELISA (peptide) LOD : 1:16000 |
| Limitations | This TRAF1 antibody is available for research use only. |

| kDa 200 – | Western blot testing of U937 cell lysate with TRAF1 antibody at 0.5ug/ml. Predicted |
|---------------|---|
| 116 – 97 – | molecular weight: ~46 kDa, routinely observed at ~50 kDa. |
| 66 | |
| 44 – | |
| 31 – | |
| 22 – | |
| 14 – | |
| 6 | |
| | |



IHC testing of FFPE human small intestine tissue with TRAF1 antibody at 2.5ug/ml. Required HIER: steamed antigen retrieval with pH6 citrate buffer; AP-staining.



IHC testing of FFPE human thymus tissue with TRAF1 antibody at 2.5ug/ml. Required HIER: steamed antigen retrieval with pH6 citrate buffer; AP-staining.

Description

Additional name(s) for this target protein: TNF receptor-associated factor 1

Application Notes

Optimal dilution of the TRAF1 antibody should be determined by the researcher.

Immunogen

Amino acids KLQSPKHAYVKDD were used as the immunogen for this TRAF1 antibody.

Storage

Aliquot and store the TRAF1 antibody at -20oC.

References (1)