


# TRAIP Antibody / TRAF-interacting protein / TRIP (RQ7536)


| Catalog No. | Formulation   | Size   |
|-------------|---|--------|
| RQ7536      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

## **Bulk quote request**


| Availability       | 1-3 business days  |
|--------------------|--|
| Species Reactivity | Human, Mouse, Rat  |
| Format             | Antigen affinity purified  |
| Clonality          | Polyclonal (rabbit origin)   |
| Isotype            | Rabbit IgG   |
| Purity             | Antigen affinity purified  |
| Buffer             | Lyophilized from 1X PBS with 2% Trehalose  |
| UniProt            | Q9BWF2   |
| Localization       | Nuclear, cytoplasmic   |
| Applications       | Western Blot : 0.5-1ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml |
| Limitations        | This TRAIP antibody is available for research use only.  |



Immunofluorescent staining of FFPE human HeLa cells with TRAIP antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human Daudi, 2) human U-251, 3) rat brain and 4) mouse brain tissue lysate with TRAIP antibody. Predicted molecular weight ~53 kDa.



Flow cytometry testing of human SiHa cells with TRAIP antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= TRAIP antibody.

## **Description**

TRAF-interacting protein is a protein that in humans is encoded by the TRAIP gene. This gene encodes a protein that contains an N-terminal RING finger motif and a putative coiled-coil domain. A similar murine protein interacts with TNFR-associated factor 1 (TRAF1), TNFR-associated factor 2 (TRAF2), and cylindromatosis. The interaction with TRAF2 inhibits TRAF2-mediated nuclear factor kappa-B, subunit 1 activation that is required for cell activation and protection against apoptosis.

### **Application Notes**

Optimal dilution of the TRAIP antibody should be determined by the researcher.

#### **Immunogen**

E. coli-derived recombinant human protein (amino acids M1-S469) was used as the immunogen for the TRAIP antibody.

#### **Storage**

After reconstitution, the TRAIP antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.