


## TRPC6 Antibody (R30847)


Catalog No.	Formulation	Size
R30847	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA, 0.025% sodium azide/thimerosal
<b>UniProt</b>	Q9Y210
<b>Applications</b>	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 0.5-1ug/ml
<b>Limitations</b>	This TRPC6 antibody is available for research use only.




Western blot testing of TRPC6 antibody and Lane 1: rat lung; 2: human 293T cell lysate.  
Predicted molecular weight: ~106 kDa.



IHC-P: TRPC6 antibody testing of human intestinal cancer tissue



IHC-P: TRPC6 antibody testing of rat brain tissue



Western blot testing of mouse lung lysate with TRPC6 antibody. Predicted molecular weight: ~106 kDa.

## Description

Transient receptor potential cation channel, subfamily C, member 6, also known as TRP6, is a transient receptor potential channel. Northern blot analysis revealed that TRPC6 is expressed primarily in placenta, lung, spleen, ovary, and small intestine. The protein is a nonselective cation channel that is activated by diacylglycerol (DAG) in a membrane-delimited fashion, independently of protein kinase C. Although TRPC3, the closest structural relative of TRPC6, is activated in the same manner, human TRPC1 and mouse *Trpc4* and *Trpc5* were unresponsive to DAG.

Immunofluorescence studies showed that most TRPC6 expression is confined to podocytes. It is also expressed in glomerular endothelial cells. Cardiac-specific overexpression of the protein in transgenic mice resulted in heightened sensitivity to stress, a propensity for lethal cardiac growth and heart failure, and an increase in Nfat-dependent expression of beta-myosin heavy chain (MYH7), a marker for pathologic hypertrophy. Studies of *Trpc6*<sup>-/-</sup> mice showed that it has a role in regulation of smooth muscle tone in blood vessels and lung.

## Application Notes

The stated application concentrations are suggested starting amounts. Titration of the TRPC6 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

An amino acid sequence from the middle region of human TRPC6 (HDYFCKCND CNQKQKHD) was used as the immunogen for this TRPC6 antibody.

## Storage

After reconstitution, the TRPC6 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

