


VCAM1 Antibody [clone VCAM1/843] (V2923)


Catalog No.	Formulation	Size
V2923-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2923-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2923SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2923IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	VCAM1/843
Purity	Protein G affinity chromatography
UniProt	P19320
Localization	Cytoplasmic, membrane
Applications	Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT
Limitations	This VCAM1 antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human tonsil stained with VCAM1 antibody (VCAM1/843).



SDS-PAGE analysis of purified, BSA-free VCAM1 antibody (clone VCAM1/843) as confirmation of integrity and purity.

Description

Recognizes a protein of 110kDa, identified as CD106 (also known as vascular cell adhesion molecule-1 (VCAM-1), L1CAM and INCAM-100). It is a member of the Ig superfamily of adhesion molecules and is expressed at high levels on cytokine stimulated vascular endothelial cells, and at minimal levels on un-stimulated endothelial cells. It is also present on follicular and inter-follicular dendritic cells of lymph nodes, myoblasts, and some macrophages. VCAM1 serves as a ligand for leukocyte integrin (VLA-4 or CD49d/CD29) and mediates cell adhesion of leukocytes to activated endothelium. It plays a role in various immunological and inflammatory responses.

Application Notes

Optimal dilution of the VCAM1 antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 min.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for the VCAM1 antibody.

Storage

Store the VCAM1 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).