



# VDAC1 Antibody (F49743)

Catalog No.	Formulation	Size
F49743-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F49743-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml


## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Predicted Reactivity	Mouse, Rat, Bovine, Rabbit
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Purified
UniProt	P21796
Applications	Western Blot : 1:1000 IHC (Paraffin) : 1:50-1:100 Flow Cytometry : 1:10-1:50
Limitations	This VDAC1 antibody is available for research use only.





IHC analysis of FFPE human hepatocarcinoma stained with VDAC1 antibody



VDAC1 antibody flow cytometry analysis of HL-60 cells (green) compared to a <a href=../search\_result.php?search\_txt=n1001>negative control</a> (blue). FITC-conjugated goat-anti-rabbit secondary Ab was used for the analysis.

### **Description**

VDAC1 forms a channel through the mitochondrial outer membrane and also the plasma membrane. The channel at the outer mitochondrial membrane allows diffusion of small hydrophilic molecules; in the plasma membrane it is involved in cell volume regulation and apoptosis. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV. The open state has a weak anion selectivity whereas the closed state is cation-selective. The protein may participate in the formation of the permeability transition pore complex (PTPC) responsible for the release of mitochondrial products that triggers apoptosis.

#### **Application Notes**

Titration of the VDAC1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

A portion of amino acids 95-124 from the human protein was used as the immunogen for this VDAC1 antibody.

#### **Storage**

Aliquot the VDAC1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.