

## VEGF Antibody / VEGFA (R30265)


| Catalog No. | Formulation   | Size   |
|-------------|---|--------|
| R30265      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |




Citations (2)

[Bulk quote request](#)


|                           |   |
|---------------------------|---|
| <b>Availability</b>       | 1-3 business days   |
| <b>Species Reactivity</b> | Human, Mouse, Rat   |
| <b>Format</b>             | Antigen affinity purified   |
| <b>Clonality</b>          | Polyclonal (rabbit origin)  |
| <b>Isotype</b>            | Rabbit IgG  |
| <b>Purity</b>             | Antigen affinity  |
| <b>Buffer</b>             | Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide       |
| <b>UniProt</b>            | P15692  |
| <b>Localization</b>       | Cytoplasmic, cell surface and extracellular (secreted)              |
| <b>Applications</b>       | Western Blot : 0.5-1ug/ml<br>Immunohistochemistry (FFPE) : 2-5ug/ml |
| <b>Limitations</b>        | This VEGF antibody is available for research use only.              |



IHC staining of FFPE human lung cancer tissue with TCP1 alpha antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human liver cancer tissue with TCP1 alpha antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human HeLa, 2) human MCF7, 3) rat kidney and 4) mouse kidney tissue lysate with VEGF antibody. Expected molecular weight 19~22 kDa (monomer) and 38~44 kDa (dimer). Banding may appear larger than predicted due to glycosylation.

## Description

VEGF, a homodimeric glycoprotein of relative molecular mass 45,000, is the only mitogen that specifically acts on endothelial cells. It may be a major regulator of tumor angiogenesis in vivo. Vascular endothelial growth factor is a mitogen primarily for vascular endothelial cells. It is, however, structurally related to platelet-derived growth factor. VEGF shares homology with the PDGF A chain and B chain, including conservation of all 8 cysteines found in PDGFA and PDGFB. VEGF gene contains 8 exons. Vascular endothelial growth factor (VEGF) induces remodeling and enhances TH2-mediated sensitization and inflammation in the lung. VEGF regulates haematopoietic stem cell survival by an internal autocrine loop mechanism. Vascular endothelial growth factor (VEGF) stimulates neurogenesis in vitro and in vivo.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the VEGF antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

An amino acid sequence from the N-terminus of human VEGFA (VDIFQEYPDEIEYIFK) was used as the immunogen for this VEGF antibody (100% homologous in human, mouse and rat).

## Storage

After reconstitution, the VEGF antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.