


von Willebrand Factor Antibody [clone WFA52-2] (V7082)


| Catalog No. | Formulation | Size |
|----------------|--|--------|
| V7082-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V7082-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V7082SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Bulk quote request

| Species Reactivity | Human |
|--------------------|---|
| Format | Purified |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG1, kappa |
| Clone Name | WFA52-2 |
| Purity | Protein G purified |
| Gene ID | 7450 |
| Localization | Cytoplasmic |
| Applications | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Limitations | This von Willebrand Factor antibody is available for research use only. |



IHC testing of FFPE human pancreas and von Willebrand Factor antibody (clone WFA52-2). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.



IHC staining of human bone marrow with von Willebrand Factor antibody (clone WFA52-2). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.

Description

Application Notes

Variations in protocols, secondaries and substrates may require the von Willebrand Factor antibody to be titered for optimal performance.

Immunogen

A recombinant human protein fragment was used as the immunogen for this von Willebrand Factor antibody.

Storage

Store the von Willebrand Factor antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).