


VPS4A Antibody (RQ6019)


Catalog No.	Formulation	Size
RQ6019	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	Q9UN37
Localization	Cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry : 1-2ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This VPS4A antibody is available for research use only.




IHC staining of FFPE human lung cancer with VPS4A antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human breast cancer with VPS4A antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) K562, 2) Raji, 3) HEK293, 4) rat brain and 5) mouse testis lysate with VPS4A antibody. Predicted molecular weight ~49 kDa.



Flow cytometry testing of human A431 cells with VPS4A antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= VPS4A antibody.

Description

Vacuolar protein sorting-associated protein 4A is a protein that in humans is encoded by the VPS4A gene. It is mapped to 16q22.1. The protein encoded by this gene is a member of the AAA protein family (ATPases associated with diverse cellular activities), and is the homolog of the yeast Vps4 protein. In humans, two paralogs of the yeast protein have been identified. The former share a high degree of aa sequence similarity with each other, and also with yeast Vps4 and mouse Skd1 proteins. The mouse Skd1 (suppressor of K⁺ transport defect 1) has been shown to be really an yeast Vps4 ortholog. Functional studies indicate that both human paralogs associate with the endosomal compartments, and are involved in intracellular protein trafficking, similar to Vps4 protein in yeast. The gene encoding this paralog has been mapped to chromosome 16; the gene for the other resides on chromosome 18.

Application Notes

Optimal dilution of the VPS4A antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids M1-K71) was used as the immunogen for the VPS4A antibody.

Storage

After reconstitution, the VPS4A antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

