


Zebrafish Csf1r Antibody / M-csfr (RZ1237)


| Catalog No. | Formulation | Size |
|-------------|---|--------|
| RZ1237 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

Bulk quote request


| Availability | 2-3 weeks |
|--------------------|---|
| Species Reactivity | Zebrafish |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Antigen affinity chromatography |
| Buffer | Lyophilized from 1X PBS with 2% Trehalose |
| UniProt | Q9I8N6 |
| Localization | Cell membrane |
| Applications | Immunohistochemistry (FFPE) : 2-5ug/ml |
| Limitations | This Zebrafish Csf1r antibody is available for research use only. |



IHC staining of zebrafish Csf1r protein using Zebrafish Csf1r antibody, HRP-labeled secondary and DAB substrate. M-Csf1r was detected in a paraffin-embedded section of zebrafish pancreas tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Csf1r protein using Zebrafish Csf1r antibody, HRP-labeled secondary and DAB substrate. Csf1r was detected in a paraffin-embedded section of zebrafish kidney tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Csf1r protein using Zebrafish Csf1r antibody, HRP-labeled secondary and DAB substrate. Csf1r was detected in a paraffin-embedded section of zebrafish spinal cord tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.