

Zebrafish Insulin like growth factor 1 Antibody / Igf1 (RZ1230)


Catalog No.	Formulation	Size
RZ1230	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	2-3 weeks
Species Reactivity	Zebrafish
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity chromatography
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q90VV9
Applications	Immunohistochemistry (FFPE) : 2-5ug/ml
Limitations	This Zebrafish Insulin like growth factor 1 antibody is available for research use only.




IHC staining of zebrafish Insulin like growth factor 1 protein using Zebrafish Insulin like growth factor 1 antibody, HRP-labeled secondary and DAB substrate. Insulin like growth factor 1 was detected in a paraffin-embedded section of zebrafish pancreas tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of zebrafish Insulin like growth factor 1 protein using Zebrafish Insulin like growth factor 1 antibody, HRP-labeled secondary and DAB substrate. Insulin like growth factor 1 was detected in a paraffin-embedded section of zebrafish brain tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Insulin like growth factor 1 protein using Zebrafish Insulin like growth factor 1 antibody, HRP-labeled secondary and DAB substrate. Insulin like growth factor 1 was detected in a paraffin-embedded section of zebrafish colon tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Insulin like growth factor 1 protein using Zebrafish Insulin like growth factor 1 antibody, HRP-labeled secondary and DAB substrate. Insulin like growth factor 1 was detected in a paraffin-embedded section of zebrafish ovary tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Insulin like growth factor 1 protein using Zebrafish Insulin like growth factor 1 antibody, HRP-labeled secondary and DAB substrate. Insulin like growth factor 1 was detected in a paraffin-embedded section of zebrafish muscle tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.