


Zebrafish Pa2g4 Antibody / Pa2g4a / Pa2g4b (RZ1260)


Catalog No.	Formulation	Size
RZ1260	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	2-3 weeks
Species Reactivity	Zebrafish
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity chromatography
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q8AW82, Q6PHD8
Localization	Cytoplasmic, Nuclear (human similarity)
Applications	Immunohistochemistry (FFPE) : 2-5ug/ml
Limitations	This Zebrafish Pa2g4 antibody is available for research use only.




IHC staining of zebrafish Pa2g4 protein using Zebrafish Pa2g4 antibody, HRP-labeled secondary and DAB substrate. Pa2g4 was detected in a paraffin-embedded section of zebrafish muscle tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Pa2g4 protein using Zebrafish Pa2g4 antibody, HRP-labeled secondary and DAB substrate. Pa2g4 was detected in a paraffin-embedded section of zebrafish heart tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Pa2g4 protein using Zebrafish Pa2g4 antibody, HRP-labeled secondary and DAB substrate. Pa2g4 was detected in a paraffin-embedded section of zebrafish liver tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Pa2g4 protein using Zebrafish Pa2g4 antibody, HRP-labeled secondary and DAB substrate. Pa2g4 was detected in a paraffin-embedded section of zebrafish brain tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.