


Zebrafish Pafah1b1 Antibody / Pafah1b1a / Pafah1b1b / Lissencephaly-1 (RZ1262)


Catalog No.	Formulation	Size
RZ1262	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)


Availability	2-3 weeks
Species Reactivity	Zebrafish
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity chromatography
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q7T394, Q803D2
Localization	Cytoplasm
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml
Limitations	This Zebrafish Pafah1b1 antibody is available for research use only.




IHC staining of FFPE zebrafish brain tissue with Pafah1b1 antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




Western blot analysis of Pafah1b1 protein using Zebrafish Pafah1b1 antibody and 1) whole female zebrafish tissue lysates and 2) whole male zebrafish tissue lysates. Predicted molecular weight ~47 kDa.




IHC staining of FFPE zebrafish heart tissue with Pafah1b1 antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE zebrafish colon tissue with Pafah1b1 antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE zebrafish muscle tissue with Pafah1b1 antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE zebrafish pancreas tissue with Pafah1b1 antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.