


## Zebrafish Psmc5 Antibody / 26S proteasome regulatory subunit 8 (RZ1290)


| Catalog No. | Formulation                                           | Size   |
|-------------|-------------------------------------------------------|--------|
| RZ1290      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

**Bulk quote request**


|                           |                                                                   |
|---------------------------|-------------------------------------------------------------------|
| <b>Availability</b>       | 2-3 weeks                                                         |
| <b>Species Reactivity</b> | Zebrafish                                                         |
| <b>Format</b>             | Antigen affinity purified                                         |
| <b>Clonality</b>          | Polyclonal (rabbit origin)                                        |
| <b>Isotype</b>            | Rabbit Ig                                                         |
| <b>Purity</b>             | Antigen affinity chromatography                                   |
| <b>Buffer</b>             | Lyophilized from 1X PBS with 2% Trehalose                         |
| <b>UniProt</b>            | Q6AZC1                                                            |
| <b>Localization</b>       | Cytoplasmic, Nuclear                                              |
| <b>Applications</b>       | Immunohistochemistry (FFPE) : 2-5ug/ml                            |
| <b>Limitations</b>        | This Zebrafish Psmc5 antibody is available for research use only. |




IHC staining of zebrafish Psmc5 protein using Zebrafish Psmc5 antibody, HRP-labeled secondary and DAB substrate. Psmc5 was detected in a paraffin-embedded section of zebrafish brain tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of zebrafish Psmc5 protein using Zebrafish Psmc5 antibody, HRP-labeled secondary and DAB substrate. Psmc5 was detected in a paraffin-embedded section of zebrafish eye tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of zebrafish Psmc5 protein using Zebrafish Psmc5 antibody, HRP-labeled secondary and DAB substrate. Psmc5 was detected in a paraffin-embedded section of zebrafish spinal tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of zebrafish Psmc5 protein using Zebrafish Psmc5 antibody, HRP-labeled secondary and DAB substrate. Psmc5 was detected in a paraffin-embedded section of zebrafish kidney tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish PSMC5 protein using Zebrafish PSMC5 antibody, HRP-labeled secondary and DAB substrate. PSMC5 was detected in a paraffin-embedded section of zebrafish heart tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Psmc5 protein using Zebrafish Psmc5 antibody, HRP-labeled secondary and DAB substrate. Psmc5 was detected in a paraffin-embedded section of zebrafish muscle tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Psmc5 protein using Zebrafish Psmc5 antibody, HRP-labeled secondary and DAB substrate. Psmc5 was detected in a paraffin-embedded section of zebrafish bone tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.