


Zebrafish Sf3b3 Antibody / Splicing factor 3b subunit 3 (RZ1308)


Catalog No.	Formulation	Size
RZ1308	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	2-3 weeks
Species Reactivity	Zebrafish
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity chromatography
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q1LVE8
Localization	Nuclear
Applications	Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence : 5ug/ml
Limitations	This Zebrafish Sf3b3 antibody is available for research use only.




Immunofluorescent staining of Sf3b3 protein using Zebrafish Sf3b3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of zebrafish Sf3b3 protein using Zebrafish Sf3b3 antibody, HRP-labeled secondary and DAB substrate. Sf3b3 was detected in a paraffin-embedded section of zebrafish brain tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Sf3b3 protein using Zebrafish Sf3b3 antibody, HRP-labeled secondary and DAB substrate. Sf3b3 was detected in a paraffin-embedded section of zebrafish heart tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Sf3b3 protein using Zebrafish Sf3b3 antibody, HRP-labeled secondary and DAB substrate. Sf3b3 was detected in a paraffin-embedded section of zebrafish muscle tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of zebrafish Sf3b3 protein using Zebrafish Sf3b3 antibody, HRP-labeled secondary and DAB substrate. Sf3b3 was detected in a paraffin-embedded section of zebrafish kidney tissue. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.