


Zebrafish Twist Antibody / Twist1a / Twist1b / Twist2 (RZ1322)


Catalog No.	Formulation	Size
RZ1322	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	2-3 weeks
Species Reactivity	Zebrafish
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity chromatography
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q9PTE3, Q568M7, Q5XJI8
Localization	Nuclear
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml
Limitations	This Zebrafish Twist antibody is available for research use only.




IHC staining of FFPE zebrafish brain tissue with Twist antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




Western blot analysis of Twist1a/1b/2 protein using Zebrafish Twist1 antibody and 1) zebrafish head tissue lysates and 2) whole male zebrafish lysates. Predicted molecular weight ~19 kDa.



IHC staining of FFPE zebrafish spinal cord tissue with Twist antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE zebrafish muscle tissue with Twist antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE zebrafish kidney tissue with Twist antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.